



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

International Recommendations for Electrocardiographic Interpretation in Athletes / Journal of the American College of Cardiology

Sharma, Sanjay ; Drezner, Jonathan A ; Baggish, Aaron ; Papadakis, Michael ; Wilson, Mathew G ; et al ; Schmied, Christian M

Abstract: Sudden cardiac death (SCD) is the leading cause of mortality in athletes during sport. A variety of mostly hereditary, structural, or electrical cardiac disorders are associated with SCD in young athletes, the majority of which can be identified or suggested by abnormalities on a resting 12-lead electrocardiogram (ECG). Whether used for diagnostic or screening purposes, physicians responsible for the cardiovascular care of athletes should be knowledgeable and competent in ECG interpretation in athletes. However, in most countries a shortage of physician expertise limits wider application of the ECG in the care of the athlete. A critical need exists for physician education in modern ECG interpretation that distinguishes normal physiological adaptations in athletes from distinctly abnormal findings suggestive of underlying pathology. Since the original 2010 European Society of Cardiology recommendations for ECG interpretation in athletes, ECG standards have evolved quickly over the last decade; pushed by a growing body of scientific data that both tests proposed criteria sets and establishes new evidence to guide refinements. On February 26-27, 2015, an international group of experts in sports cardiology, inherited cardiac disease, and sports medicine convened in Seattle, Washington, to update contemporary standards for ECG interpretation in athletes. The objective of the meeting was to define and revise ECG interpretation standards based on new and emerging research and to develop a clear guide to the proper evaluation of ECG abnormalities in athletes. This statement represents an international consensus for ECG interpretation in athletes and provides expert opinion-based recommendations linking specific ECG abnormalities and the secondary evaluation for conditions associated with SCD.

DOI: <https://doi.org/10.1016/j.jacc.2017.01.015>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-206521>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Sharma, Sanjay; Drezner, Jonathan A; Baggish, Aaron; Papadakis, Michael; Wilson, Mathew G; et al; Schmied, Christian M (2017). International Recommendations for Electrocardiographic Interpretation in

Athletes / Journal of the American College of Cardiology. Journal of the American College of Cardiology, 69(8):1057-1075.
DOI: <https://doi.org/10.1016/j.jacc.2017.01.015>

CURRENT OPINION

International Recommendations for Electrocardiographic Interpretation in Athletes



Sanjay Sharma, MD,^{a,*} Jonathan A. Drezner, MD,^{b,*} Aaron Baggish, MD,^c Michael Papadakis, MD,^a Mathew G. Wilson, PhD,^d Jordan M. Prutkin, MD, MHS,^e Andre La Gerche, MD, PhD,^f Michael J. Ackerman, MD, PhD,^g Mats Borjesson, MD, PhD,^h Jack C. Salerno, MD,ⁱ Irfan M. Asif, MD,^j David S. Owens, MD, MS,^e Eugene H. Chung, MD, MS,^k Michael S. Emery, MD,^l Victor F. Froelicher, MD,^m Hein Heidbuchel, MD, PhD,^{n,o} Carmen Adamuz, MD, PhD,^d Chad A. Asplund, MD,^p Gordon Cohen, MD,^q Kimberly G. Harmon, MD,^b Joseph C. Marek, MD,^r Silvana Molossi, MD,^s Josef Niebauer, MD, PhD,^t Hank F. Pelto, MD,^b Marco V. Perez, MD,^u Nathan R. Riding, PhD,^d Tess Saarel, MD,^v Christian M. Schmied, MD,^w David M. Shipon, MD,^x Ricardo Stein, MD, ScD,^y Victoria L. Vetter, MD, MPH,^z Antonio Pelliccia, MD,^{aa} Domenico Corrado, MD, PhD^{bb}

ABSTRACT

Sudden cardiac death (SCD) is the leading cause of mortality in athletes during sport. A variety of mostly hereditary, structural, or electrical cardiac disorders are associated with SCD in young athletes, the majority of which can be identified or suggested by abnormalities on a resting 12-lead electrocardiogram (ECG). Whether used for diagnostic or screening purposes, physicians responsible for the cardiovascular care of athletes should be knowledgeable and competent in ECG interpretation in athletes. However, in most countries a shortage of physician expertise limits wider application of the ECG in the care of the athlete. A critical need exists for physician education in modern ECG interpretation that distinguishes normal physiological adaptations in athletes from distinctly abnormal findings suggestive of underlying pathology. Since the original 2010 European Society of Cardiology recommendations for ECG interpretation in athletes, ECG standards have evolved quickly over the last decade; pushed by a growing body of scientific data that both tests proposed criteria sets and establishes new evidence to guide refinements. On February 26-27, 2015, an international group of experts in sports cardiology, inherited cardiac disease, and sports medicine convened in Seattle, Washington, to update contemporary standards for ECG interpretation in athletes. The objective of the meeting was to define and revise ECG interpretation standards based on new and emerging research and to develop a clear guide to the proper evaluation of ECG abnormalities in athletes. This statement represents an international consensus for ECG interpretation in athletes and provides expert opinion-based recommendations linking specific ECG abnormalities and the secondary evaluation for conditions associated with SCD. (J Am Coll Cardiol 2017;69:1057-75) © 2017 The Authors. Published by Elsevier Inc. on behalf of American College of Cardiology Foundation. All rights reserved.

This article has been co-published in the *European Heart Journal* and the *Journal of the American College of Cardiology*. An extended version of this article has also been jointly published in the *British Journal of Sports Medicine*.

From the ^aCardiology Clinical and Academic Group, St George's University of London, United Kingdom; ^bDepartment of Family Medicine, University of Washington, Seattle, Washington; ^cDivision of Cardiology, Massachusetts General Hospital, Boston, Massachusetts; ^dDepartment of Sports Medicine, ASPETAR, Qatar Orthopaedic and Sports Medicine Hospital, Qatar; ^eDivision of Cardiology, University of Washington, Seattle, Washington; ^fDepartment of Cardiology, Baker IDI Heart and Diabetes Institute, Melbourne, Australia; ^gDepartment of Cardiovascular Diseases, Pediatric and Adolescent Medicine, and Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, Minnesota; ^hDepartment of Neuroscience and Physiology, Sahlgrenska University Hospital/Ostra Sahlgrenska Academy, Goteborg, Sweden; ⁱDepartment of Pediatrics, University of Washington, Seattle, Washington; ^jDepartment of Family Medicine, University of South Carolina, Greenville, South Carolina; ^kDivision of Cardiology, University of North Carolina School of Medicine, Chapel Hill, North Carolina; ^lCenter of Cardiovascular Care in Athletics, Indiana University School of Medicine, Indianapolis, Indiana; ^mDepartment of Medicine, Stanford University, Stanford, California; ⁿDepartment of Cardiology, Arrhythmology Hasselt University, Belgium; ^oDepartment of Cardiology, Antwerp, Belgium; ^pGeorgia

ABBREVIATIONS AND ACRONYMS

ARVC = arrhythmogenic right ventricular cardiomyopathy
AV = atrioventricular
BrS = Brugada syndrome
DCM = dilated cardiomyopathy
ECG = electrocardiogram
HCM = hypertrophic cardiomyopathy
IVCD = intra-ventricular conduction delay
LBBB = left bundle branch block
LGE = late gadolinium enhancement
LQTS = long QT syndrome
LVH = left ventricular hypertrophy
LVNC = left ventricular noncompaction
MRI = magnetic resonance imaging
RBBB = right bundle branch block
RVH = right ventricular hypertrophy
SAECG = signal average electrocardiogram
SCD = sudden cardiac death
SVT = supraventricular tachycardia
TWI = T-wave inversion
VF = ventricular fibrillation
VPC = ventricular premature contractions
WPW = Wolf-Parkinson-White

INTRODUCTION

Cardiovascular-related sudden death is the leading cause of mortality in athletes during sport and exercise (1-3). The majority of disorders associated with an increased risk of sudden cardiac death (SCD) are suggested or identified by abnormalities on a resting 12-lead ECG. Whether used for the evaluation of cardiovascular-related symptoms, a family history of inheritable cardiac disease or premature SCD, or for screening of asymptomatic athletes, ECG interpretation is an essential skill for all physicians involved in the cardiovascular care of athletes.

THE 2015 SUMMIT ON ECG INTERPRETATION IN ATHLETES. Over the last decade, ECG interpretation standards have undergone several modifications to improve the accuracy of detecting potentially life threatening cardiac conditions in young athletes while also limiting false positive results (4-15). In February 2015, an international group of experts convened in Seattle, Washington, to update contemporary recommendations for ECG interpretation in asymptomatic athletes age 12 to 35 years. The goals of the summit meeting were to: 1) update ECG interpretation standards based on new and emerging research; and 2) develop a clear guide to the appropriate evaluation of ECG abnormalities for conditions associated with SCD in athletes. In the presence of cardiac symptoms or a family history of inherited cardiovascular disease or premature SCD, a normal ECG should not preclude further assessment.

This document provides the most updated evidence-based recommendations developed with

thoughtful attention to balance sensitivity and specificity, while maintaining a clear and practical checklist of findings to guide ECG interpretation for physicians and the appropriate evaluation of ECG abnormalities. A summary of the consensus recommendations is presented in [Figure 1](#) and [Tables 1 and 2](#).

LIMITATIONS. While ECG increases the ability to detect underlying cardiovascular conditions associated with SCD, ECG as a diagnostic tool has limitations in both sensitivity and specificity. Specifically, ECG is unable to detect anomalous coronary arteries, premature coronary atherosclerosis, and aortopathies. In some instances patients with cardiomyopathies, particularly arrhythmogenic right ventricular cardiomyopathy (ARVC), may also reveal a normal ECG. Thus, an ECG will not detect all conditions predisposing to SCD. Furthermore, inter-observer variability among physicians remains a major concern (16-18), despite studies demonstrating that using standardized criteria improves interpretation accuracy (19,20).

NORMAL ECG FINDINGS IN ATHLETES


PHYSIOLOGICAL CARDIAC ADAPTATIONS TO REGULAR EXERCISE. Regular and long-term participation in intensive exercise (minimum of 4 h per week) is associated with unique electrical manifestations that reflect enlarged cardiac chamber size and increased vagal tone. These ECG findings in athletes are considered normal, physiological adaptations to regular exercise and do not require further evaluation ([Figure 1](#), [Table 1](#)).

LEFT AND RIGHT VENTRICULAR HYPERTROPHY. The presence of isolated QRS voltage criterion for left ventricular hypertrophy (LVH) ([Figure 2](#)) does not correlate with pathology in athletes and is present in isolation (without other associated ECG abnormalities) in <2% of patients with hypertrophic

Southern University, Statesboro, Georgia; ⁴Division of Pediatric Cardiothoracic Surgery, University of California San Francisco School of Medicine, San Francisco, California; ⁵Advocate Heart Institute, Normal, Illinois; ⁶Division of Pediatric Cardiology, Baylor College of Medicine, Houston, Texas; ⁷University Institute of Sports Medicine, Paracelsus Medical University, Austria; ⁸Center for Inherited Cardiovascular Disease, Stanford University, Stanford, California; ⁹Pediatric Cardiology, Cleveland Clinic, Cleveland, Ohio; ¹⁰University of Herzzentrum, Zurich, Switzerland; ¹¹Heart Center of Philadelphia, Jefferson University Hospitals, Philadelphia, Pennsylvania; ¹²Department of Cardiology, Hospital de Clinicas de Porte Alegre, Brazil; ¹³The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania; ¹⁴Institute of Sports Medicine and Science, Rome, Italy; and the ¹⁵Department of Cardiac, Thoracic and Vascular Sciences, University of Padua Medical School, Italy. Dr. Sharma has received grants from Cardiac Risk in the Young; and grants from British Heart Foundation, outside the submitted work. Dr. Papadakis has received grants from Cardiac Risk in the Young outside the submitted work. Dr. Ackerman has received personal fees from Boston Scientific, Gilead Sciences, Invitae, Medtronic, St. Jude Medical, and other from Transgenomic, outside the submitted work; in addition, Dr. Ackerman has a patent QT and T Wave Analytics pending. Dr. Froelicher has received other from insightinc, from null, outside the submitted work; this is an entity that manufactures and develops ECG software and devices for screening purposes. Dr. Heidbuchel has received other from Biotronik, during the conduct of the study; and personal fees from Biotronik, Pfizer/BMS, Daiichi-Sankyo, Bayer, Boehringer-Ingelheim, Cardione, outside the submitted work. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

*The first 2 authors contributed equally to the study.

FIGURE 1 International Consensus Standards for Electrocardiographic Interpretation in Athletes



cardiomyopathy (HCM) (21-27). Conversely, pathological LVH is commonly associated with additional ECG features such as T-wave inversion (TWI) in the inferior and lateral leads, ST-segment depression, and pathological Q waves (28,29). Therefore, the isolated presence of high QRS voltages fulfilling voltage criterion for LVH in the absence of other ECG or clinical markers suggestive of pathology are considered part of normal and training-related ECG changes in athletes and does not require further evaluation.

Voltage criterion for right ventricular hypertrophy (RVH) is also common in athletes with up to 13% of athletes fulfilling the Sokolow-Lyon index (30,31). QRS voltages for RVH, when present in isolation, do not correlate with underlying pathology in athletes (31). Similar to voltage criteria for LVH, isolated QRS voltage for RVH is part of the normal spectrum of ECG findings in athletes and does not require further evaluation.

EARLY REPOLARIZATION. Early repolarization is defined as elevation of the QRS-ST junction (J-point) by ≥ 0.1 mV often associated with a late QRS slurring or notching (J-wave) affecting the inferior and/or lateral leads (32-34). Early repolarization is common in healthy populations (2% to 44%) and is more prevalent

in athletes, young individuals, males, and black ethnicity (32,35-39). Early repolarization consisting of J-point elevation with concave ST-segment elevation and a peaked TWI (Figure 2) is present in up to 45% of Caucasian athletes and 63% to 91% of black athletes of African-Caribbean descent (hereto referred to as 'black' athletes) (22,30).

Some studies on survivors of cardiac arrest and patients with primary ventricular fibrillation (VF) have suggested an association between early repolarization and the risk of VF (33,40). Although further studies are warranted to fully elucidate the mechanisms and prognostic implications of early repolarization in competitive athletes, to date there are no data to support an association between inferior early repolarization and SCD in athletes. Based on current evidence, all patterns of early repolarization, when present in isolation and without clinical markers of pathology, should be considered benign variants in athletes (41).

REPOLARIZATION FINDINGS IN BLACK ATHLETES. Ethnicity is a major determinant of cardiac adaptation to exercise with more than two-thirds of black athletes exhibiting repolarization changes (29,30,42,43). Black athletes also commonly demonstrate a repolarization

TABLE 1 International Consensus Standards for Electrocardiographic Interpretation in Athletes: Definitions of ECG Criteria

Abnormal ECG findings in athletes	
These ECG findings are unrelated to regular training or expected physiologic adaptation to exercise, may suggest the presence of pathologic cardiovascular disease, and require further diagnostic investigation.	
ECG abnormality	Definition
T wave inversion	≥1 mm in depth in two or more contiguous leads; excludes leads aVR, III, and V ₁
• Anterior	• V ₂ -V ₄ <ul style="list-style-type: none"> - excludes: black athletes with J-point elevation and convex ST-segment elevation followed by TWI in V₂-V₄; athletes age <16 with TWI in V₁-V₃; and biphasic T waves in only V₃
• Lateral	• I and AVL, V ₅ and/or V ₆ (only one lead of TWI required in V ₅ or V ₆)
• Inferolateral	• II and aVF, V ₅ -V ₆ , I and AVL
• Inferior	• II and aVF
ST-segment depression	≥0.5 mm in depth in two or more contiguous leads
Pathologic Q waves	Q/R ratio ≥0.25 or ≥40 ms in duration in two or more leads (excluding III and aVR)
Complete left bundle branch block	QRS ≥120 ms, predominantly negative QRS complex in lead V ₁ (QS or rS), and upright notched or slurred R wave in leads I and V ₆
Profound nonspecific intra-ventricular conduction delay	Any QRS duration ≥140 ms
Epsilon wave	Distinct low amplitude signal (small positive deflection or notch) between the end of the QRS complex and onset of the T-wave in leads V ₁ -V ₃
Ventricular pre-excitation	PR interval <120 ms with a delta wave (slurred upstroke in the QRS complex) and wide QRS (≥120 ms)
Prolonged QT interval*	QTc ≥470 ms (male) QTc ≥480 ms (female) QTc ≥500 ms (marked QT prolongation)
Brugada Type 1 pattern	Coved pattern: initial ST-segment elevation ≥2 mm (high take-off) with downsloping ST-segment elevation followed by a negative symmetric T-wave in ≥ 1 leads in V ₁ -V ₃
Profound sinus bradycardia	<30 beats/min or sinus pauses ≥3 s
Profound 1° AV block	≥400 ms
Mobitz Type II 2° AV block	Intermittently non-conducted P waves with a fixed PR interval
3° AV block	Complete heart block
Atrial tachyarrhythmias	Supraventricular tachycardia, atrial fibrillation, atrial flutter
PVC	≥2 PVCs per 10 s tracing
Ventricular arrhythmias	Couplets, triplets, and non-sustained ventricular tachycardia
Borderline ECG findings in athletes	
These ECG findings in isolation likely do not represent pathologic cardiovascular disease in athletes, but the presence of two or more borderline findings may warrant additional investigation until further data become available.	
ECG abnormality	Definition
Left axis deviation	-30° to -90°
Left atrial enlargement	Prolonged P wave duration of >120 ms in leads I or II with negative portion of the P-wave ≥1 mm in depth and ≥40 ms in duration in lead V ₁
Right axis deviation	>120°
Right atrial enlargement	P-wave ≥2.5 mm in II, III, or aVF
Complete right bundle branch block	rSR' pattern in lead V ₁ and a S wave wider than R wave in lead V ₆ with QRS duration ≥120 ms

Continued on the next page

variant consisting of J-point elevation and convex ST-segment elevation in the anterior leads (V₁ to V₄) followed by TWI (**Figures 3, 4B, and 4C**) which is regarded as a normal variant and should not result in further investigation, in the absence of other clinical or ECG features of cardiomyopathy (30,42-44).

CONSIDERATIONS IN ATHLETES AGE 12 TO 16 YEARS: THE 'JUVENILE' ELECTROCARDIOGRAM PATTERN.

TWI confined to the anterior precordial leads may be considered a normal age-related pattern in adolescent athletes up to the age of 16 years old. The term 'juvenile' ECG pattern is used to denote TWI or a biphasic T-wave beyond lead V₂ in adolescents who have not reached physical maturity and is present in

10%-15% of white, adolescent athletes aged 12 years old but only in 2.5% of white athletes age 14 to 15 years (**Figure 4A**) (22,45,46). Anterior TWI that extends beyond lead V₂ is rare (0.1%) in white athletes age ≥16 years or younger athletes who have completed puberty (22,45). Based on current evidence, TWI in the anterior leads (V₁ to V₃) in adolescent athletes <16 years of age should not prompt further evaluation in the absence of symptoms, signs, or a family history of cardiac disease.

PHYSIOLOGICAL ARRHYTHMIAS IN ATHLETES. Common consequences of increased vagal tone include sinus bradycardia and sinus arrhythmia (22,47-49). Other, less common markers of increased vagal tone are

TABLE 1 Continued

Normal ECG findings in athletes

These training-related ECG alterations are physiologic adaptations to regular exercise, considered normal variants in athletes, and do not require further evaluation in asymptomatic athletes with no significant family history.

Normal ECG finding	Definition
Increased QRS voltage	Isolated QRS voltage criteria for left (SV1 + RV5 or RV6 >3.5 mV) or right ventricular hypertrophy (RV1 + SV5 or SV6 >1.1 mV)
Incomplete RBBB	rSR' pattern in lead V ₁ and a qRS pattern in lead V ₆ with QRS duration <120 ms
Early repolarization	J-point elevation, ST-segment elevation, J waves, or terminal QRS slurring in the inferior and/or lateral leads
Black athlete repolarization variant	J-point elevation and convex ('domed') ST-segment elevation followed by T-wave inversion in leads V ₁ –V ₄ in black athletes
Juvenile T-wave pattern	T-wave inversion V ₁ –V ₃ in athletes age <16 yrs
Sinus bradycardia	≥30 beats/min
Sinus arrhythmia	Heart rate variation with respiration: rate increases during inspiration and decreases during expiration
Ectopic atrial rhythm	P waves are a different morphology compared with the sinus P-wave, such as negative P waves in the inferior leads ('low atrial rhythm')
Junctional escape rhythm	QRS rate is faster than the resting P-wave or sinus rate and typically <100 beats/min with narrow QRS complex unless the baseline QRS is conducted with aberrancy
1° AV block	PR interval 200–400 ms
Mobitz Type I (Wenckebach) 2° AV block	PR interval progressively lengthens until there is a non-conducted P-wave with no QRS complex; the first PR interval after the dropped beat is shorter than the last conducted PR interval

*The QT interval corrected for heart rate is ideally measured using Bazett's formula with heart rates between 60 and 90 beats/min; preferably performed manually in lead II or V₅ using the 'teach-the-tangent method' to avoid inclusion of a U-wave (please see text for more details). Consider repeating the ECG after mild aerobic activity for a heart rate <50 beats/min, or repeating the ECG after a longer resting period for a heart rate >100 beats/min, if the QTc value is borderline or abnormal.

AV = atrioventricular block; ECG = electrocardiogram; PVC = premature ventricular contraction; RBBB = right bundle branch block.

junctional or ectopic atrial rhythms, first degree atrio-ventricular (AV) block, and Mobitz Type I second degree AV block (Wenckebach phenomenon) (22,47,50). In the absence of symptoms, heart rates ≥ 30 beats/min are considered normal in highly trained athletes. Sinus rhythm should resume and bradycardia should resolve with the onset of physical activity.

BORDERLINE ELECTROCARDIOGRAM FINDINGS IN ATHLETES

Recent data suggest that some ECG findings previously categorized as abnormal may represent normal variants or the result of physiological cardiac remodeling in athletes and do not usually represent pathological cardiac disease. These ECG findings have been categorized as 'borderline' findings in athletes (Figure 1, Table 1).

AXIS DEVIATION AND VOLTAGE CRITERIA FOR ATRIAL ENLARGEMENT. Axis deviation and voltage criteria for atrial enlargement account for >40% of abnormal ECG patterns in athletes but do not correlate with cardiac pathology (51). In a large study of 2,533 athletes age 14 to 35 years old and 9,997 controls of similar age, echocardiographic evaluation of the 579 athletes and controls with isolated axis deviation or voltage criteria for atrial enlargement failed to identify any major structural or functional abnormalities (51).

COMPLETE RIGHT BUNDLE BRANCH BLOCK. Although incomplete right bundle branch block (RBBB) is

common in young athletes, the significance of complete RBBB is less certain. Complete RBBB is detected in approximately 1% of the general population and large datasets in young adult athletes reveal a prevalence of 0.5% to 2.5% (12,52–54). In a study of 510 U.S. collegiate athletes, RBBB was reported in 2.5% and compared with athletes with normal QRS complexes or incomplete RBBB, athletes with complete RBBB exhibited larger right ventricular dimensions and a lower right ventricular ejection fraction but preserved fractional area change (55). None of the athletes with complete RBBB or incomplete RBBB was found to have pathological structural cardiac disease. These patterns among trained athletes could represent a spectrum of structural and physiological cardiac remodeling characterized by RV dilation with resultant QRS prolongation and a relative reduction in the RV systolic function at rest (55).

Based on the aforementioned considerations, left axis deviation, left atrial enlargement, right axis deviation and right atrial enlargement and complete RBBB are considered borderline variants in athletes. The presence of any one of these findings in isolation or with other recognized physiological electrical patterns of athletic training does not warrant further assessment in asymptomatic athletes without a family history of premature cardiac disease or SCD. Conversely, the presence of more than one of these borderline findings places the athlete in the abnormal category warranting additional investigation.

TABLE 2 Evaluation of Electrocardiographic Abnormalities

ECG Abnormality	Potential Cardiac Disease*	Recommended Evaluation†	Considerations
T-wave inversion in the lateral or inferolateral leads	HCM DCM LVNC ARVC (with predominant LV involvement) Myocarditis	Echocardiography CMR Exercise ECG test Minimum 24 h ECG monitor	Lateral or inferolateral T wave inversion is common in primary myocardial disease. CMR should be a routine diagnostic test for this ECG phenotype and is superior to echocardiography for detecting apical HCM, LVH localized to the free lateral wall, ARVC with predominant left ventricular involvement, and myocarditis. If CMR is not available, echocardiography with contrast should be considered as an alternative investigation for apical HCM in patients with deep T wave inversion in leads V ₅ -V ₆ . Consider family evaluation if available and genetic screening. Annual follow-up testing is recommended throughout athletic career in athletes with normal results.
T-wave inversion isolated to the inferior leads	HCM DCM LVNC Myocarditis	Echocardiography	Consider CMR based on echocardiography findings or clinical suspicion.
T-wave inversion in the anterior leads‡	ARVC DCM	Echocardiography CMR Exercise ECG test Minimum 24 h ECG monitor SAECG	The extent of investigations may vary based on clinical suspicion for ARVC and results from initial testing.
ST-segment depression	HCM DCM LVNC ARVC Myocarditis	Echocardiography	Consider CMR and additional testing based on echocardiography findings or clinical suspicion.
Pathologic Q waves	HCM DCM LVNC Myocarditis Prior MI	Echocardiography CAD risk factor assessment Repeat ECG for septal (V ₁ -V ₂) QS pattern; above investigations recommended if septal Q waves are persistent	Consider CMR (with perfusion study if available) based on echocardiography findings or clinical suspicion. In the absence of CMR, consider exercise stress testing, dobutamine stress echocardiogram, or a myocardial perfusion scan for evaluation of coronary artery disease in athletes with suspicion of prior MI or multiple risk factors for CAD.
Complete left bundle branch block	DCM HCM LVNC Sarcoidosis Myocarditis	Echocardiography CMR (with stress perfusion study)§	A comprehensive cardiac evaluation to rule out myocardial disease should be considered.
Profound nonspecific intraventricular conduction delay ≥140 ms	DCM HCM LVNC	Echocardiography	Consider additional testing based on echocardiography findings or clinical suspicion.
Epsilon wave	ARVC	Echocardiography CMR Exercise ECG test Minimum 24 h ECG monitor SAECG	An epsilon wave in leads V ₁ -V ₃ is a highly specific ECG marker and a major diagnostic criterion for ARVC.
Multiple premature ventricular contractions	HCM DCM LVNC ARVC Myocarditis Sarcoidosis	Echocardiography 24 h ECG monitor Exercise ECG test	If >2,000 PVC's or non-sustained ventricular tachycardia are present on initial testing, comprehensive cardiac testing inclusive of CMR is warranted to investigate for myocardial disease. Consider signal averaged ECG (SAECG).
Ventricular pre-excitation	WPW	Exercise ECG test Echocardiography	Abrupt cessation of the delta wave (pre-excitation) on exercise ECG denotes a low risk pathway. EP study for risk assessment should be considered if a low risk accessory pathway cannot be confirmed by non-invasive testing. Consider EP study for moderate to high intensity sports.
Prolonged QTc	LQTS	Repeat resting ECG on separate day Review for QT prolonging medication Acquire ECG of 1st degree relatives if possible	Consider exercise ECG test, laboratory (electrolyte) screening, family screening and genetic testing when clinical suspicion is high. Consider direct referral to a heart rhythm specialist or sports cardiologist for a QTc ≥500 ms.
Brugada Type 1 pattern	Brugada syndrome	Referral to cardiologist or heart rhythm specialist	Consider high precordial lead ECG with leads V ₁ and V ₂ in 2nd intercostal space or sodium channel blockade if Brugada pattern is indeterminate. Consider genetic testing and family screening.
Profound sinus bradycardia <30 beats/min	Myocardial or electrical disease	Repeat ECG after mild aerobic activity	Consider additional testing based on clinical suspicion.
Profound 1° atrioventricular block ≥400 ms	Myocardial or electrical disease	Repeat ECG after mild aerobic activity Exercise ECG test	Consider additional testing based on clinical suspicion.

Continued on the next page

TABLE 2 Continued

ECG Abnormality	Potential Cardiac Disease*	Recommended Evaluation†	Considerations
Advanced 2° or 3° atrioventricular block	Myocardial or electrical disease	Echocardiography Minimum 24 h ECG monitor Exercise ECG test	Consider laboratory screening and CMR based on echocardiography findings.
Atrial tachyarrhythmias	Myocardial or electrical disease	Echocardiography Minimum 24 h ECG monitor Exercise ECG test	Consider CMR or EP study based on clinical suspicion.
Ventricular arrhythmias	Myocardial or electrical disease	Echocardiography CMR Minimum 24 h ECG monitor Exercise ECG test	A comprehensive cardiac evaluation to rule out myocardial disease and primary electrical disease should be considered.
Two or more borderline ECG findings	Myocardial disease	Echocardiography	Consider additional testing based on clinical suspicion.

*This list of disorders for each ECG abnormality represents the primary cardiac disorders of concern and is not intended to be exhaustive. †Initial evaluation of ECG abnormalities should be performed under the direction of a cardiologist. Additional testing will be guided by initial findings and clinical suspicion based on the presence of symptoms or a family history of inherited cardiac disease or SCD. ‡Excludes black athlete repolarization variant and juvenile pattern in adolescents < 16 years. §CT coronary angiography if stress perfusion with CMR is unavailable. ||Includes couplets, triplets, accelerated ventricular rhythm, and non-sustained ventricular tachycardia.

ARVC = arrhythmogenic right ventricular cardiomyopathy; CAD = coronary artery disease; CMR = cardiovascular magnetic resonance; DCM = dilated cardiomyopathy; ECG = electrocardiogram; EP = electrophysiological; HCM = hypertrophic cardiomyopathy; LQTS = long QT syndrome; LVNC = left ventricular noncompaction; MI = myocardial infarction; PVC = premature ventricular contraction; SAEKG = signal averaged electrocardiogram; WPPW = Wolff Parkinson White syndrome.

ABNORMAL ELECTROCARDIOGRAM FINDINGS IN ATHLETES

The abnormal findings defined in this section are not recognized features of athletic training and always require further assessment to exclude the presence of intrinsic cardiac disease (Figure 1, Tables 1 and 2). Temporary restriction from athletic activity should be considered for athletes with abnormal ECGs of uncertain clinical significance until secondary investigations are completed.

ABNORMAL T-WAVE INVERSION. TWI ≥ 1 mm in depth in two or more contiguous leads (excluding leads aVR, III, and V₁) in an anterior, lateral, inferolateral, or inferior territory is abnormal and should prompt further evaluation for underlying structural heart disease (Tables 1 and 2). Normal exceptions include TWI confined to leads V₁–V₄ in black athletes when preceded by J point and/or ST-segment elevation, and TWI in leads V₁–V₃ in athletes aged < 16 years.

CLINICAL CONSIDERATIONS. The relationship between abnormal TWI and several forms of structural heart disease is well documented (56). TWI in the inferior or lateral leads is common in HCM (56–59). Whereas TWI in the right precordial leads (V₁ to V₃) or beyond in the absence of a complete RBBB is common in ARVC (Figure 4D) (60,61).

There are no data relating to the significance of flat or biphasic T-waves in athletes but similar to TWI, this panel would recommend further evaluation of biphasic T-waves where the negative portion is ≥ 1 mm in depth in ≥ 2 leads.


Evaluation. Lateral or inferolateral T-wave inversion. There is mounting evidence that TWI in the lateral or

inferolateral leads is associated with the presence of quiescent cardiomyopathy in a considerable proportion of athletes (30,62–64). Recommendations for the evaluation of abnormal TWI and other clinical considerations are presented in Table 2.

TWI affecting the lateral leads (V₅ to V₆, I and aVL) (Figure 4E) should prompt a comprehensive investigation to exclude cardiomyopathy. If echocardiography is not diagnostic, cardiac magnetic resonance imaging (MRI) with gadolinium should be utilized. Cardiac MRI provides superior assessment of myocardial hypertrophy, especially the left ventricular apex and the lateral free wall and may also demonstrate late gadolinium enhancement (LGE), a non-specific marker suggesting myocardial fibrosis. If cardiac MRI is not available, echocardiography with contrast should be considered. Exercise ECG testing and Holter monitoring also should be considered in the evaluation of lateral or inferolateral TWI, especially for athletes with ‘grey zone’ hypertrophy (males with maximal LV wall thickness 13 to 16 mm) without LGE, where the diagnosis of HCM remains uncertain. In such cases, the presence of ventricular tachycardia during exercise or Holter may support HCM and is also useful in risk stratification (65).

For athletes with lateral or inferolateral TWI, regular follow-up with serial cardiac imaging is necessary even when the initial evaluation is normal, in order to monitor for the development of a cardiomyopathy phenotype (62,63).

Anterior T-wave inversion. Anterior TWI is a normal variant in asymptomatic adolescent athletes age <16 years, in black athletes when preceded by J-point elevation and convex ST-segment elevation, and in some endurance athletes (66,67). However, anterior

FIGURE 2 Early Repolarization Changes in an Athlete

Electrocardiogram of a 29-year-old male asymptomatic soccer player showing sinus bradycardia (44 beats/min), early repolarization in I, II, aVF, V₂ to V₆ (arrows), voltage criterion for left ventricular hypertrophy (S-V₁ + R-V₅ > 35 mm), and tall, peaked T waves (circles). These are common, training related findings in athletes and do not require more evaluation.

TWI in leads V₁ to V₂/V₃ also is a recognized pattern in patients with ARVC and rarely HCM.


There are discrepancies among existing guidelines relating to the extent of anterior TWI inversion before considering further investigations (5,6,14,29). A large study of over 14,000 white adults age 16 to 35 years old, including over 2,500 athletes showed that anterior TWI had a prevalence of 2.3% (68). Anterior TWI was more common in females and athletes and was confined to leads V₁ to V₂ in almost all individuals, and only exceeded beyond V₂ in 1% of females and 0.2% of males (68). None of the individuals with anterior TWI were diagnosed with a cardiomyopathy following comprehensive investigation indicating that this particular ECG pattern is non-specific in low-risk populations. Based on this report, it is justifiable to only investigate non-black athletes with anterior TWI beyond V₂ in the absence of other clinical or electrical features of ARVC.

Specific information about the J-point and preceding ST-segment may help differentiate between physiological adaptation and cardiomyopathy in athletes with anterior TWI affecting leads V₃ and/or V₄. A recent study comparing anterior TWI in a series of black and white healthy athletes, and patients with HCM and ARVC, showed that in athletes with anterior

TWI, the combination of J-point elevation ≥ 1 mm and TWI confined to leads V₁ to V₄ excluded either cardiomyopathy with 100% negative predictive value, regardless of ethnicity (66). Conversely, anterior TWI associated with minimal or absent J-point elevation (<1 mm) could reflect a cardiomyopathy (66). These data require duplication in larger studies but may prove useful in the assessment of a small proportion of white endurance athletes who exhibit anterior TWI and in athletes of black/mixed ethnicity (69).

In most non-black athletes age ≥ 16 years, anterior TWI beyond lead V₂ should prompt further evaluation given the potential overlap with ARVC. In these athletes, concurrent findings of J-point elevation, ST-segment elevation, or biphasic T waves more likely represents athlete's heart, while the absence of J-point elevation or a coexistent depressed ST-segment is more concerning for ARVC (Figure 5) (66). Other ECG findings suggestive of ARVC include low limb lead voltages, prolonged S wave upstroke, ventricular ectopy with LBBB morphology, and epsilon waves (61). A combination of tests is needed to diagnose ARVC including echocardiography, cardiac MRI, Holter monitoring, exercise ECG test, and signal averaged ECG.

FIGURE 3 Anterior Repolarization Changes in a Black Athlete



Electrocardiogram from a black athlete demonstrating voltage criterion for left ventricular hypertrophy, J-point elevation, and convex ('domed') ST-segment elevation followed by T-wave inversion in V₁ to V₄ (circles). This is a normal repolarization pattern in black athletes.

Inferior T-wave inversion. The significance of TWI confined to the inferior leads is unknown. However, this finding cannot be attributed to physiological remodeling and thus warrants further investigation with, at minimum, an echocardiogram. Cardiac MRI should be considered based on the echocardiographic findings or clinical suspicion.

ST-SEGMENT DEPRESSION. While ST-segment depression is common among patients with cardiomyopathy, it is not a feature of athletic training (28,59,70,71). ST-segment depression (relative to the isoelectric PR segment) in excess of 0.05 mV (0.5 mm) in two or more leads should be considered an abnormal finding requiring definitive evaluation for underlying structural heart disease.


Evaluation. Echocardiography is the minimum evaluation for athletes with ST-segment depression to investigate for underlying cardiomyopathy. Cardiac MRI should be considered based on the echocardiographic findings or clinical suspicion.

PATHOLOGICAL Q WAVES. Several pathological disorders including HCM, ARVC, infiltrative myocardial diseases, accessory pathways and transmural myocardial infarction can lead to the development of

exaggerated (deep or wide) Q waves or unexpected Q waves in atypical leads (28,56). Pathological Q waves also may be a result of lead misplacement. In particular, a pseudo-septal infarct pattern with pathological Q waves in leads V₁ to V₂ is commonly due to high-lead placement relative to cardiac position (72).

Pathological Q waves have been reported in approximately 1% to 2% of all athletes, and may be higher in males and black athletes (29,73). For asymptomatic athletes, pathological Q waves were previously defined as >3 mm in depth or >40 ms in duration in two or more leads (except III and aVR) (6,10). In practice, however, this criterion is a common source of false positive ECG results as trained athletes with physiological LVH and thin adolescent athletes may have increased precordial voltages and deep lateral or inferior Q waves.

The use of a Q/R ratio overcomes some of these issues by normalizing Q wave depth to the degree of preceding R-wave voltage. Case control analyses of athletes and HCM patients suggest that this will decrease the false positive rate without compromising sensitivity for the detection of cardiomyopathy (29,74). Thus, this consensus panel has modified the definition for pathological Q waves

FIGURE 4 Normal and Abnormal Patterns of T-Wave Inversion

(A) Anterior T-wave inversion in V₁ to V₃ in a 12-year-old asymptomatic athlete without a family history of sudden cardiac death considered a normal 'juvenile' pattern. (B) T-wave inversion in V₁ to V₄ in a 17-year-old asymptomatic mixed race (Middle-Eastern/black) athlete without a family history of sudden cardiac death. This is a normal repolarization pattern in black athletes. (C) Biphasic T-wave inversion in V₃ in a 31-year-old asymptomatic black athlete without a family history of sudden cardiac death. Anterior biphasic T waves are considered normal in adolescents <16 years old and in adults when found in a single lead, most commonly V₃. (D) Abnormal T-wave inversion in V₁ to V₆ in an adult symptomatic former soccer player with genetically confirmed arrhythmogenic right ventricular cardiomyopathy and a positive family history of sudden cardiac death (brother died at 26 years of age). (E) Inferolateral T-wave inversion in leads I, II, III, aVF, V₂ to V₆, and ST-segment depression in leads II, aVF, V₄ to V₆ in a 31-year-old asymptomatic professional soccer referee. These markedly abnormal findings require a comprehensive evaluation to exclude cardiomyopathy. (F) An electrocardiogram demonstrating the Type 1 Brugada pattern with high take-off ST-segment elevation ≥ 2 mm with downsloping ST-segment elevation followed by a negative symmetric T wave in V₁ to V₂.


in athletes as a Q/R ratio ≥ 0.25 or ≥ 40 ms in duration in two or more contiguous leads (except III and aVR).

Evaluation. An ECG with abnormal Q waves should be carefully examined for the possibility of an accessory pathway. If the pathological Q waves are isolated to leads V₁ to V₂, the ECG should be repeated, including re-placing the ECG leads to ensure proper positioning. Persistence of pathological Q waves in two or more contiguous leads warrants further investigation with echocardiography to exclude cardiomyopathy. If the echocardiogram is normal and there are no other

concerning clinical findings or ECG abnormalities, no additional testing is generally necessary. However, if there is a high index of clinical suspicion, additional evaluation with cardiac MRI should be considered. In athlete's age ≥ 30 years with suspicion of prior myocardial infarction or risk factors for coronary artery disease (CAD), stress testing may be warranted.

COMPLETE LEFT BUNDLE BRANCH BLOCK. LBBB is found in <1 in 1,000 athletes but is common in patients with cardiomyopathy and ischemic heart

FIGURE 5 Examples of Physiological and Pathological T-Wave Inversion



(A) T-wave inversion in V₁ to V₄ preceded by J-point elevation and convex 'domed' ST-segment elevation (**green circles**). This should not be confused with pathological T-wave inversion (**B**) which demonstrates T-wave inversion in V₁ to V₆ with absent J-point elevation and a downsloping ST-segment (**red circles**).

disease (9,28,59,75,76). Thus, complete LBBB always should be considered an abnormal finding and requires a comprehensive evaluation to rule out a pathological cardiac disorder.

Evaluation. Athletes with complete LBBB require a thorough investigation for myocardial disease including echocardiography and a cardiac MRI with perfusion study.

PROFOUND NON-SPECIFIC INTRA-VENTRICULAR CONDUCTION DELAY. Epidemiological studies of nonspecific intra-ventricular conduction delay (IVCD) in the general population have shown an increased risk of cardiovascular death and have been documented among patients with cardiomyopathy (77,78). The significance of non-specific IVCD with normal QRS morphology in healthy, asymptomatic athletes is uncertain (79). The physiology underlying IVCD in athletes remains incompletely understood but likely includes some combination of neurally mediated conduction fiber slowing and increased myocardial mass. In patients with LVH, left ventricular mass seems to be closely related to QRS duration (80).

While the exact cut-off to trigger more investigation in athletes with a nonspecific IVCD remains unclear, this panel recommends that marked

nonspecific IVCD ≥ 140 ms in athletes, regardless of QRS morphology, is abnormal and should prompt further evaluation.

Evaluation. In asymptomatic athletes with profound non-specific IVCD, an echocardiogram is recommended to evaluate for myocardial disease. Other testing may be indicated depending on echocardiographic findings or clinical suspicion.

VENTRICULAR PRE-EXCITATION. Ventricular pre-excitation occurs when an accessory pathway bypasses the AV node resulting in abnormal conduction to the ventricle (pre-excitation) with shortening of the PR interval and widening of the QRS. This is evident on the ECG as the Wolf-Parkinson-White (WPW) pattern defined as a PR interval < 120 ms, the presence of a delta wave (slurring of the initial QRS), and a QRS duration > 120 ms (81). The WPW pattern occurs in up to 1 in 250 athletes (9,12,52,82). The presence of an accessory pathway can predispose an athlete to sudden death because rapid conduction of atrial fibrillation across the accessory pathway can result in VF.

Evaluation. A short PR interval in isolation without a widened QRS or delta wave in an asymptomatic athlete should not be considered for further assessment. The WPW pattern warrants further assessment of the

refractory period of the accessory pathway. Non-invasive risk stratification begins with an exercise stress test, where abrupt, complete loss of pre-excitation at higher heart rates suggests a low-risk accessory pathway (83,84). An echocardiogram also should be considered due to the association of WPW with Ebstein's anomaly and cardiomyopathy. Intermittent pre-excitation during sinus rhythm on a resting ECG is also consistent with a low-risk pathway and may obviate the need for an exercise test (85). If non-invasive testing cannot confirm a low-risk pathway or is inconclusive, an electrophysiological study should be considered to determine the shortest pre-excited RR interval during atrial fibrillation (83). If the shortest pre-excited RR interval is ≤ 250 ms (240 beats/min), then the accessory pathway is deemed high risk and transcatheter ablation is recommended (83,86). Some physicians may choose to subject all competitive athletes involved in moderate or high-intensity sport to electrophysiological studies irrespective of the results of the exercise test or 24 h ECG on the premise that high catecholamine concentrations during very intensive exercise may modify the refractory period of an accessory pathway in a fashion that cannot be reproduced during laboratory tests.

PROLONGED QT INTERVAL. Congenital long QT syndrome (LQTS) is a potentially lethal, genetically mediated ventricular arrhythmia syndrome with the hallmark electrocardiographic feature of QT prolongation. LQTS is estimated to affect 1 in 2,000 individuals, and this may be underestimated given the subpopulation of so-called 'normal QT interval' or 'concealed' LQTS (87). Autopsy negative sudden unexplained death represents 25% to 40% of sudden unexpected deaths in persons under age 40 years (3,88-90). In such cases, cardiac ion channelopathies have been implicated by post-mortem genetic testing as the probable cause in up to 25% to 40% of cases (91-94).

Calculating the corrected QT interval. Accurate measurement and manual confirmation of the computer derived QT interval corrected for heart rate (QTc) is critical as the accuracy of computer generated QTc values is about 90% to 95%. Studies have suggested the ability of cardiologists to accurately measure the QTc is suboptimal (95). However, accurate assessment of the QTc can be achieved by adhering to the following six principles (96):


1. Use Bazett's heart rate correction formula ($QTc = QT/\sqrt{RR}$; note the RR interval is measured in seconds) as population-based QTc distributions most frequently use Bazett-derived QTc values (97).
2. Bazett's formula underestimates the QTc at heart rates <50 beats/min, and overestimates the QTc at heart rates >90 beats/min. Accordingly, for a heart rate <50 beats/min, a repeat ECG after mild aerobic activity is recommended to achieve a heart rate closer to 60 beats/min. For heart rates >90 beats/min, a repeat ECG after additional resting time may help achieve a lower heart rate.
3. If sinus arrhythmia is present with beat to beat variation in heart rate, an average QT interval and average RR interval should be used.
4. Leads II and V_5 usually provide the best delineation of the T-wave.
5. Low amplitude U waves, which are common in the anterior precordial leads, should not be included in the QT calculation. The 'Teach-the-Tangent' or 'Avoid-the-Tail' method to delineate the end of the T-wave should be followed (Figure 6) (96).
6. The morphology of the T-wave, not just the length of the QT interval, also can suggest the presence of LQTS (98). For instance, a notched T-wave in the lateral precordial leads where the amplitude of the second portion of the T-wave following the notch is greater than the first portion of the T-wave may represent LQT-2 even in the absence of overt QT prolongation.

The easiest and most efficient way to confirm the computer-derived QTc is to examine lead II and/or V_5 and determine if the manually measured QT interval matches the computer's QT measurement. If there is concordance within about 10 ms, one can trust that the computer can derive accurately an average RR interval and complete the Bazett's calculation. If, however, the manually measured QT interval is >10 ms different than the computer's QT measurement, an average RR interval should be determined and the QTc recalculated using the Bazett's formula.

Corrected QT cut-offs. Given the overlap between QTc distributions in population-derived cohorts of healthy individuals compared with patients with genetically confirmed LQTS, the QTc cut-off value compelling further evaluation must be chosen carefully to balance the frequency of abnormal results and the positive predictive value for LQTS.

Recent consensus statements on ECG interpretation in athletes have recommended that male athletes with a $QTc \geq 470$ ms and female athletes with a $QTc \geq 480$ ms undergo further evaluation for LQTS to better balance false positive and false negative findings (6,10). These cut-off values are around the 99th percentile and consistent with thresholds defined by the American Heart Association and

FIGURE 6 This Figure Illustrates the 'Teach-the-Tangent' or 'Avoid-the-Tail' Method for Manual Measurement of the QT Interval



A straight line is drawn on the downslope of the T-wave to the point of intersection with the isoelectric line. The U-wave is not included in the measurement.

American College of Cardiology (99). This consensus group also recommends QTc values of ≥ 470 ms in males and ≥ 480 ms in females to define the threshold of QT prolongation that warrants further assessment in asymptomatic athletes.

Short QT interval. The precise cut-off and clinical significance of a short QT interval in athletes is unknown. Data from over 18,000 asymptomatic young British individuals found that the prevalence of a QTc < 320 ms is 0.1%; suggesting an abnormal cut-off value of < 320 ms is pragmatic (100). However, over a mean follow up period of 5.3 years, none of the individuals with a short QT < 320 ms experienced any adverse events, syncope, or sudden death (100). Based on the rarity of this finding and absence of data to suggest long-term morbidity in asymptomatic athletes, this panel recommends that a short QT interval only be investigated in the context of concerning clinical markers.


Evaluation. It is critical that an athlete with a single prolonged QTc reading not be obligated a diagnosis of LQTS, but rather that these cut-off values trigger the need for additional evaluation. The importance of additional evaluation but not a premature diagnosis of LQTS was demonstrated in a study of 2,000 elite athletes in which 7 (0.4%) had a prolonged QTc (range

460 to 570 ms) (101). A QTc of < 500 ms in the absence of symptoms or familial disease was unlikely to represent LQTS. In contrast, a QTc ≥ 500 ms was highly suggestive of LQTS as all three athletes with a QTc value of > 500 ms exhibited one of paradoxical prolongation of the QTc during exercise, a confirmatory genetic mutation, or prolonged QTc in a first-degree relative (101).

A personal history of syncope or seizures and a family history of exertional syncope, 'epilepsy', postpartum-timed syncope/seizure, unexplained motor vehicle accidents, unexplained drowning, and premature, unexplained sudden death < 50 years of age should be reviewed. If the personal/family history is positive, the athlete should be referred to an electrophysiologist for further evaluation. If the personal/family history is negative, a repeat ECG should be obtained (ideally on a different day). If the follow-up ECG is below the QTc cut-off values, then no additional evaluation is needed and the athlete should be reassured.

If the repeat ECG still exceeds the QTc cut-off values, then a screening ECG of the athlete's first degree relatives (parents and siblings) should be considered and the athlete should be referred to an electrophysiologist for the possibility of newly

FIGURE 7 Differentiation Between the Brugada ECG Pattern From Early Repolarization (J Point Elevation and Convex ST Segment) in Athletes



Brugada Type 1 electrocardiogram (**left**) should be distinguished from early repolarization with 'convex' ST-segment elevation in a trained athlete (**right**). Vertical lines mark the J-point (STJ) and the point 80 ms after the J-point (ST80), where the amplitudes of the ST-segment elevation are calculated. The 'downsloping' ST-segment elevation in Brugada pattern is characterized by a STJ/ST80 ratio >1 . Early repolarization patterns in an athlete show an initial 'upsloping' ST-segment elevation with STJ/ST80 ratio <1 .

discovered LQTS. Reversible, extrinsic factors, such as electrolyte abnormalities (hypokalaemia) or the presence of QT prolonging medications, must also be evaluated. If an athlete's ECG shows a QTc ≥ 500 ms and no reversible causes are identified, then the athlete should be referred immediately to an electrophysiologist as the probability of LQTS and future adverse events has increased (102). The Schwartz-Moss scoring system, electrocardiographic features, stress ECG, provocative testing, and genetic testing may be needed to clarify the diagnosis and should be performed and interpreted by a cardiologist familiar with the disease (103-106).

BRUGADA TYPE 1 PATTERN. Brugada syndrome (BrS) is an inherited primary electrical disease which predisposes to ventricular tachyarrhythmias and sudden death during states of enhanced vagal tone. It is characterized by the distinctive Brugada ECG pattern which consists of a coved rSr' pattern, ST-segment elevation ≥ 2 mm, and inversion of the terminal portion of the T-wave in leads V₁, V₂, and V₃ (Figure 4F). Although three types were described, only the Type 1 Brugada pattern is now considered diagnostic (107-109).

The coved ST-segment elevation in Type 1 Brugada pattern results in a broad r' and should be distinguishable from the upsloping ST-segment elevation of early repolarization in an athlete. In this regards, the 'Corrado index' measures the ST-segment elevation at the start of the ST-segment/J-point (STJ) and 80 ms after the start of the ST-segment (ST80)

(110). In Type 1 Brugada pattern, the downsloping ST-segment will have a STJ/ST80 ratio >1 , while the initial upsloping of the ST-segment found in early repolarization patterns in an athlete will produce an STJ/ST80 ratio <1 (Figure 7).

Evaluation. The Type 1 Brugada ECG pattern should be investigated regardless of symptoms. If the pattern is unclear, confirm correct lead placement, repeat the ECG if necessary, and perform a high precordial lead ECG with V₁ and V₂ placed in the 2nd or 3rd intercostal space. If the Type 1 pattern is seen on a high-precordial lead ECG, then referral to an electrophysiologist is indicated. Consideration should be given to potential accentuating factors for a Brugada-like ECG pattern, such as hyperkalaemia, fever, medications with sodium ion channel blocking properties, and lead placement.

PROFOUND SINUS BRADYCARDIA OR FIRST DEGREE ATRIOVENTRICULAR BLOCK. Sinus bradycardia and moderate prolongation of the PR interval (200-399 ms) are recognized features of athletic conditioning. Although a resting heart rate ≤ 30 beats/min or a PR interval ≥ 400 ms may be normal in a well-trained athlete, it should prompt further evaluation for cardiac conduction disease.

Evaluation. Evaluation of profound sinus bradycardia or a markedly increased PR interval should include assessing the chronotropic response to mild aerobic activity, such as running on the spot or climbing stairs. Exercise testing is useful in this

situation to provide an objective measure of the PR interval and heart rate response to aerobic activity. If the heart rate increases appropriately and the PR interval normalizes, and the athlete is asymptomatic, no further testing is necessary. Conversely, further evaluation should be performed if the heart rate does not increase or the PR interval does not shorten appropriately on exertion, the athlete experiences pre-syncope/syncope, or in athletes with a family history of cardiac disease or sudden death. Depending on the clinical scenario, an echocardiogram or ambulatory ECG monitor may be indicated.

HIGH GRADE ATRIOVENTRICULAR BLOCK. Mobitz Type II second degree AV block and third degree (complete) AV block are abnormal findings in athletes. Complete heart block can be confused with AV dissociation without block; a situation where the junctional pacemaker is faster than the sinus node, leading to more QRS complexes than P waves. Intermittent ventricular capture by sinus P waves (resulting in an irregular ventricular response) excludes complete AV block. AV dissociation without block is the expression of autonomic mismatch between AV and sinus nodal modulation, but is not pathological. Like all other functional disturbances, a small exercise load with repeat ECG recording will show resolution of the ECG findings in AV dissociation.

Evaluation. If Mobitz II AV block or complete AV block is detected, further evaluation includes an echocardiogram, ambulatory ECG monitor, and exercise ECG test. Based on these results, laboratory testing and cardiac MRI may be considered. Referral to an electrophysiologist is essential.

MULTIPLE PREMATURE VENTRICULAR CONTRACTIONS. Multiple (≥ 2) premature ventricular contractions (PVCs) are uncommon and present in $<1\%$ of 12-lead ECGs in athletes (9,12). Although multiple PVCs are usually benign, their presence may be the hallmark of underlying heart disease (111,112). PVCs originating from the right ventricular outflow tract (LBBB and inferior axis origin) are considered particularly benign when associated with a normal ECG, however this PVC morphology can also be present in patients with early ARVC particularly when the QRS exceeds 160 ms (113). Therefore, the finding of ≥ 2 PVCs on an ECG should prompt more extensive evaluation to exclude underlying structural heart disease.

Evaluation. The extent of evaluation for ≥ 2 PVCs is controversial and excluding pathology may be difficult. At a minimum, an ambulatory Holter monitor, echocardiogram, and exercise stress test should be performed. The availability of modern small, leadless ambulatory recorders allows for

longer electrocardiographic monitoring, including during training and competition, to exclude complex ventricular arrhythmias. If the Holter and echocardiogram are normal and the PVCs suppress with exercise, no further evaluation is recommended for an asymptomatic athlete. A previous study has shown that among athletes with $\geq 2,000$ PVCs per 24 h, up to 30% were found to have underlying structural heart disease, compared with 3% and 0% in those with $<2,000$ and <100 PVCs per day, respectively (112). Therefore, in athletes with $\geq 2,000$ PVCs per 24 h or with episodes of non-sustained ventricular tachycardia, or with an increasing burden of ectopy during an incremental exercise test, additional evaluation may include contrast-enhanced cardiac MRI and more invasive electrophysiology study (114,115). Although some studies have suggested that regression of the PVC burden with detraining indicates a good prognosis, other studies have not confirmed this (116–118). Thus, detraining as a diagnostic or therapeutic measure is not recommended.

ATRIAL TACHYARRHYTHMIAS. Sinus tachycardia is the most common atrial tachyarrhythmia but is very rarely due to intrinsic cardiac disease. Supraventricular tachycardia (SVT), atrial fibrillation, and atrial flutter are rarely seen on a resting ECG in athletes and require investigation. Atrial tachyarrhythmias are rarely life threatening but can be associated with other conditions that can lead to SCD, including LQTS, WPW, BrS, myocarditis, congenital heart disease, and the cardiomyopathies.

Evaluation. If resting sinus tachycardia >120 beats/min is seen, a repeat ECG should be considered after a period of rest as recent exercise or anxiety may be the cause. Other underlying etiologies may be sought, including fever, infection, dehydration, stimulant use, anemia, hyperthyroidism, or, rarely, underlying cardiac or pulmonary disease.

For paroxysmal SVT, a repeat ECG when not in SVT should be obtained if possible. If the Valsalva maneuver, carotid sinus massage, or the diving reflex is used to terminate the arrhythmia, a rhythm strip should be obtained which can help elucidate the mechanism of the SVT. An echocardiogram, ambulatory ECG monitor, and exercise treadmill test should be completed. Referral to an electrophysiologist may be indicated for consideration of electrophysiology study and ablation.

If atrial fibrillation or flutter is found, an echocardiogram should be completed to assess for structural heart disease and anti-coagulation considered based on standard guidelines (119). An ambulatory ECG

monitor should be used to assess if the rhythm is paroxysmal or persistent and what the ventricular rate is throughout the day. A thorough family history may elucidate an underlying genetic cause. Depending on what these results show, cardiac MRI, electrophysiology study with possible ablation, and/or genetic testing may be considered.

VENTRICULAR ARRHYTHMIAS. Ventricular couplets, triplets, and non-sustained ventricular tachycardia always require investigation as they can be a marker for underlying cardiac pathology or lead to sustained ventricular tachycardia which may cause SCD.

Evaluation. If ventricular arrhythmias are seen, the evaluation should include a thorough family history, an echocardiogram to evaluate for structural heart disease, cardiac MRI to assess for ARVC or other cardiomyopathies, ambulatory ECG monitor and exercise ECG test. Depending on these results, further evaluation may be needed including electrophysiology study or genetic testing.

CONSIDERATIONS IN ATHLETES ≥ 30 YEARS OF AGE. In athletes ≥ 30 years of age, CAD is the most common cause of SCD (89,90). In addition, older athletes may be less fit compared with 20 to 30 years ago, increasing the possibility of underlying CAD (120,121). While resting ECGs have a low sensitivity for CAD, some ECG patterns may suggest underlying CAD such as TWI, pathological Q waves, ST-segment depression, left or RBBB, abnormal R-wave progression, left anterior hemiblock, and atrial fibrillation (122-124).

Evaluation. The main role of a resting ECG in older athletes is to identify those athletes who may potentially be at high risk for CAD and warrant further testing (123,125,126). Initial testing should include an exercise stress test, resting echocardiogram, and assessment of traditional risk factors for CAD. When indicated, this evaluation may be complemented by coronary CT angiography or a functional stress test.

ELECTROCARDIOGRAM PATTERNS REQUIRING SERIAL EVALUATION. Several common heritable cardiomyopathies may present with ECG abnormalities prior to the onset of overt heart muscle pathology (62,63). Therefore, athletes with abnormal ECGs suggestive of cardiomyopathy and initially normal clinical evaluations should be followed with serial evaluation during and after their competitive athletic careers. Evaluations may be conducted annually or more frequently depending on individual circumstances. These athletes may be permitted to participate in competitive athletics without restriction contingent on longitudinal follow-up.

CONCLUSION

Accurate ECG interpretation in athletes requires adequate training and an attention to detail to distinguish physiological ECG findings from abnormal ECG findings that might indicate the presence of cardiac pathology. The international consensus standards presented on ECG interpretation and the evaluation of ECG abnormalities serve as an important foundation for improving the quality of cardiovascular care of athletes. As new scientific data become available, revision of these recommendations may be necessary to further advance the accuracy of ECG interpretation in the athletic population.

ACKNOWLEDGEMENTS The *2nd Summit on Electrocardiogram Interpretation in Athletes* was sponsored by the American Medical Society for Sports Medicine (AMSSM), the FIFA Medical Assessment and Research Center (F-MARC), and the National Collegiate Athletic Association (NCAA). Participating medical societies included the American College of Cardiology (ACC) Sports & Exercise Council, and the Section on Sports Cardiology of the European Association for Cardiovascular Prevention and Rehabilitation (EACPR), a registered branch of the European Society of Cardiology (ESC). This statement has been endorsed by the following societies: American Medical Society for Sports Medicine (AMSSM), Austrian Society of Sports Medicine and Prevention, Brazilian Society of Cardiology - Department of Exercise and Rehabilitation (SBC - DERC), British Association for Sports and Exercise Medicine (BASEM), Canadian Academy of Sport and Exercise Medicine (CASEM), European College of Sports and Exercise Physicians (ECOSEP), European Society of Cardiology (ESC) Section of Sports Cardiology, Fédération Internationale de Football Association (FIFA), German Society of Sports Medicine and Prevention, International Olympic Committee (IOC), Norwegian Association of Sports Medicine and Physical Activity (NIMF), South African Sports Medicine Association (SASMA), Spanish Society of Cardiology (SEC) Sports Cardiology Group, Sports Doctors Australia, and the Swedish Society of Exercise and Sports Medicine (SFAIM). The American College of Cardiology (ACC) affirms the value of this document (ACC supports the general principles in the document and believes it is of general benefit to its membership).

ADDRESS FOR CORRESPONDENCE: Dr. Sanjay Sharma, Cardiology Clinical and Academic Group, St George's University of London, United Kingdom. E-mail: sasharma@sgul.ac.uk.

REFERENCES

- Harmon KG, Asif IM, Klossner D, Drezner JA. Incidence of sudden cardiac death in national collegiate athletic association athletes. *Circulation* 2011;123:1594–600.
- Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980–2006. *Circulation* 2009;119:1085–92.
- Harmon KG, Asif IM, Maleszewski JJ, et al. Incidence, etiology, and comparative frequency of sudden cardiac death in NCAA Athletes: a decade in review. *Circulation* 2015;132:10–9.
- Corrado D, Biffi A, Basso C, Pelliccia A, Thiene G. 12-lead ECG in the athlete: physiological versus pathological abnormalities. *Br J Sports Med* 2009;43:669–76.
- Corrado D, Pelliccia A, Heidbuchel H, et al. Recommendations for interpretation of 12-lead electrocardiogram in the athlete. *Eur Heart J* 2010;31:243–59.
- Uberoi A, Stein R, Perez MV, et al. Interpretation of the electrocardiogram of young athletes. *Circulation* 2011;124:746–57.
- Williams ES, Owens DS, Drezner JA, Prutkin JM. Electrocardiogram interpretation in the athlete. *Herzschrittmacherther Elektrophysiol* 2012;23:65–71.
- Drezner J. Standardised criteria for ECG interpretation in athletes: a practical tool. *Br J Sports Med* 2012;46:i6–8.
- Marek J, Bufalino V, Davis J, et al. Feasibility and findings of large-scale electrocardiographic screening in young adults: data from 32,561 subjects. *Heart Rhythm* 2011;8:1555–9.
- Drezner JA, Ackerman MJ, Anderson J, et al. Electrocardiographic interpretation in athletes: the 'Seattle criteria'. *Br J Sports Med* 2013;47:122–4.
- Corrado D, Pelliccia A, Bjornstad HH, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005;26:516–24.
- Pelliccia A, Culasso F, Di Paolo FM, et al. Prevalence of abnormal electrocardiograms in a large, unselected population undergoing pre-participation cardiovascular screening. *Eur Heart J* 2007;28:2006–10.
- Drezner JA, Fischbach P, Froelicher V, et al. Normal electrocardiographic findings: recognising physiological adaptations in athletes. *Br J Sports Med* 2013;47:125–36.
- Drezner JA, Ashley E, Baggish AL, et al. Abnormal electrocardiographic findings in athletes: recognising changes suggestive of cardiomyopathy. *Br J Sports Med* 2013;47:137–52.
- Drezner JA, Ackerman MJ, Cannon BC, et al. Abnormal electrocardiographic findings in athletes: recognising changes suggestive of primary electrical disease. *Br J Sports Med* 2013;47:153–67.
- Brosnan M, La Gerche A, Kumar S, Lo W, Kalman J, Prior D. Modest agreement in ECG interpretation limits the application of ECG screening in young athletes. *Heart Rhythm* 2015;12:130–6.
- Magee C, Kazman J, Haigney M, et al. Reliability and validity of clinician ECG interpretation for athletes. *Ann Noninvasive Electrocardiol* 2014;19:319–29.
- Hill AC, Miyake CY, Grady S, Dubin AM. Accuracy of interpretation of preparticipation screening electrocardiograms. *J Pediatr* 2011;159:783–8.
- Drezner JA, Asif IM, Owens DS, et al. Accuracy of ECG interpretation in competitive athletes: the impact of using standardised ECG criteria. *Br J Sports Med* 2012;46:335–40.
- Exeter DJ, Elley CR, Fulcher ML, Lee AC, Drezner JA, Asif IM. Standardised criteria improve accuracy of ECG interpretation in competitive athletes: a randomised controlled trial. *Br J Sports Med* 2014;48:1167–71.
- Ryan MP, Cleland JG, French JA, et al. The standard electrocardiogram as a screening test for hypertrophic cardiomyopathy. *Am J Cardiol* 1995;76:689–94.
- Papadakis M, Basavarajiah S, Rawlins J, et al. Prevalence and significance of T-wave inversions in predominantly Caucasian adolescent athletes. *Eur Heart J* 2009;30:1728–35.
- Pelliccia A, Maron BJ, Culasso F, et al. Clinical significance of abnormal electrocardiographic patterns in trained athletes. *Circulation* 2000;102:278–84.
- Sohaib SM, Payne JR, Shukla R, World M, Pennell DJ, Montgomery HE. Electrocardiographic (ECG) criteria for determining left ventricular mass in young healthy men; data from the LARGE Heart study. *J Cardiovasc Magn Reson* 2009;11:2.
- Sathanandam S, Zimmerman F, Davis J, Marek J. Abstract 2484: ECG screening criteria for LVH does not correlate with diagnosis of hypertrophic cardiomyopathy. *Circulation* 2009;S647.
- Weiner RB, Hutter AM, Wang F, et al. Performance of the 2010 European Society of Cardiology criteria for ECG interpretation in the athlete. *Heart* 2011;97:1573–7.
- Calore C, Melacini P, Pelliccia A, et al. Prevalence and clinical meaning of isolated increase of QRS voltages in hypertrophic cardiomyopathy versus athlete's heart: relevance to athletic screening. *Int J Cardiol* 2013;168:4494–7.
- Lakdawala NK, Thune JJ, Maron BJ, et al. Electrocardiographic features of sarcomere mutation carriers with and without clinically overt hypertrophic cardiomyopathy. *Am J Cardiol* 2011;108:1606–13.
- Sheikh N, Papadakis M, Ghani S, et al. Comparison of electrocardiographic criteria for the detection of cardiac abnormalities in elite black and white athletes. *Circulation* 2014;129:1637–49.
- Papadakis M, Carre F, Kervio G, et al. The prevalence, distribution, and clinical outcomes of electrocardiographic repolarization patterns in male athletes of African/Afro-Caribbean origin. *Eur Heart J* 2011;32:2304–13.
- Zaidi A, Ghani S, Sheikh N, et al. Clinical significance of electrocardiographic right ventricular hypertrophy in athletes: comparison with arrhythmogenic right ventricular cardiomyopathy and pulmonary hypertension. *Eur Heart J* 2013;34:3649–56.
- Tikkanen JT, Anttonen O, Junttila MJ, et al. Long-term outcome associated with early repolarization on electrocardiography. *N Engl J Med* 2009;361:2529–37.
- Haissaguerre M, Derval N, Sacher F, et al. Sudden cardiac arrest associated with early repolarization. *N Engl J Med* 2008;358:2016–23.
- Macfarlane PW, Antzelevitch C, Haissaguerre M, et al. The early repolarization pattern: a consensus paper. *J Am Coll Cardiol* 2015;66:470–7.
- Tikkanen JT, Junttila MJ, Anttonen O, et al. Early repolarization: electrocardiographic phenotypes associated with favorable long-term outcome. *Circulation* 2011;123:2666–73.
- Uberoi A, Jain NA, Perez M, et al. Early repolarization in an ambulatory clinical population. *Circulation* 2011;124:2208–14.
- Junttila MJ, Sager SJ, Freiser M, McGonagle S, Castellanos A, Myerburg RJ. Inferolateral early repolarization in athletes. *J Interv Card Electrophysiol* 2011;31:33–8.
- Noseworthy PA, Weiner R, Kim J, et al. Early repolarization pattern in competitive athletes: clinical correlates and the effects of exercise training. *Circ Arrhythm Electrophysiol* 2011;4:432–40.
- Noseworthy PA, Tikkanen JT, Porthan K, et al. The early repolarization pattern in the general population: clinical correlates and heritability. *J Am Coll Cardiol* 2011;57:2284–9.
- Rosso R, Kogan E, Belhassen B, et al. J-point elevation in survivors of primary ventricular fibrillation and matched control subjects: incidence and clinical significance. *J Am Coll Cardiol* 2008;52:1231–8.
- Quattrini FM, Pelliccia A, Assorgi R, et al. Benign clinical significance of J-wave pattern (early repolarization) in highly trained athletes. *Heart Rhythm* 2014;11:1974–82.
- Sheikh N, Papadakis M, Carre F, et al. Cardiac adaptation to exercise in adolescent athletes of African ethnicity: an emergent elite athletic population. *Br J Sports Med* 2013;47:585–92.
- Di Paolo FM, Schmied C, Zerguini YA, et al. The athlete's heart in adolescent Africans: an electrocardiographic and echocardiographic study. *J Am Coll Cardiol* 2012;59:1029–36.
- Rawlins J, Carre F, Kervio G, et al. Ethnic differences in physiological cardiac adaptation to intense physical exercise in highly trained female athletes. *Circulation* 2010;121:1078–85.

45. Migliore F, Zorzi A, Michieli P, et al. Prevalence of cardiomyopathy in Italian asymptomatic children with electrocardiographic T-wave inversion at preparticipation screening. *Circulation* 2012;125:529-38.
46. Calo L, Sperandii F, Martino A, et al. Electrocardiographic findings in 2261 peri-pubertal athletes with or without inverted T waves at electrocardiogram. *Heart* 2015;101:193-200.
47. Sharma S, Whyte G, Elliott P, et al. Electrocardiographic changes in 1000 highly trained junior elite athletes. *Br J Sports Med* 1999;33:319-24.
48. Stein R, Medeiros CM, Rosito GA, Zimmerman LJ, Ribeiro JP. Intrinsic sinus and atrioventricular node electrophysiologic adaptations in endurance athletes. *J Am Coll Cardiol* 2002;39:1033-8.
49. Northcote RJ, Canning GP, Ballantyne D. Electrocardiographic findings in male veteran endurance athletes. *Br Heart J* 1989;61:155-60.
50. Meytes I, Kaplinsky E, Yahini JH, Hanne-Paparo N, Neufeld HN. Wenckebach A-V block: a frequent feature following heavy physical training. *Am Heart J* 1975;90:426-30.
51. Gati S, Sheikh N, Ghani S, et al. Should axis deviation or atrial enlargement be categorised as abnormal in young athletes? The athlete's electrocardiogram: time for re-appraisal of markers of pathology. *Eur Heart J* 2013;34:3641-8.
52. Fudge J, Harmon KG, Owens DS, et al. Cardiovascular screening in adolescents and young adults: a prospective study comparing the Preparticipation Physical Evaluation Monograph 4th Edition and ECG. *Br J Sports Med* 2014;48:1172-8.
53. Magalski A, McCoy M, Zabel M, et al. Cardiovascular screening with electrocardiography and echocardiography in collegiate athletes. *Am J Med* 2011;124:S11-8.
54. Baggish AL, Hutter AM Jr., Wang F, et al. Cardiovascular screening in college athletes with and without electrocardiography: A cross-sectional study. *Ann Intern Med* 2010;152:269-75.
55. Kim JH, Noseworthy PA, McCarty D, et al. Significance of electrocardiographic right bundle branch block in trained athletes. *Am J Cardiol* 2011;107:1083-9.
56. Rowin EJ, Maron BJ, Appelbaum E, et al. Significance of false negative electrocardiograms in preparticipation screening of athletes for hypertrophic cardiomyopathy. *Am J Cardiol* 2012;110:1027-32.
57. Chen X, Zhao T, Lu M, et al. The relationship between electrocardiographic changes and CMR features in asymptomatic or mildly symptomatic patients with hypertrophic cardiomyopathy. *Int J Cardiovasc Imaging* 2014;30 Suppl 1:55-63.
58. Sheikh N, Papadakis M, Schnell F, et al. Clinical profile of athletes with hypertrophic cardiomyopathy. *Circ Cardiovasc Imaging* 2015;8:e003454.
59. Bent RE, Wheeler MT, Hadley D, et al. Systematic comparison of digital electrocardiograms from healthy athletes and patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2015;65:2462-3.
60. Nasir K, Bomma C, Tandri H, et al. Electrocardiographic features of arrhythmogenic right ventricular dysplasia/cardiomyopathy according to disease severity: a need to broaden diagnostic criteria. *Circulation* 2004;110:1527-34.
61. Marcus FI, McKenna WJ, Sherrill D, et al. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria. *Circulation* 2010;121:1533-41.
62. Pelliccia A, Di Paolo FM, Quattrini FM, et al. Outcomes in athletes with marked ECG repolarization abnormalities. *N Engl J Med* 2008;358:152-61.
63. Schnell F, Riding N, O'hannon R, et al. Recognition and significance of pathological T-wave inversions in athletes. *Circulation* 2015;131:165-73.
64. Chandra N, Bastiaenen R, Papadakis M, et al. Prevalence of electrocardiographic anomalies in young individuals: relevance to a nationwide cardiac screening program. *J Am Coll Cardiol* 2014;63:2028-34.
65. Elliott PM, Anastasakis A, Borger MA, et al. 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: the Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). *Eur Heart J* 2014;35:2733-79.
66. Calore C, Zorzi A, Sheikh N, et al. Electrocardiographic anterior T-wave inversion in athletes of different ethnicities: differential diagnosis between athlete's heart and cardiomyopathy. *Eur Heart J* 2015;37:2515-27.
67. Brosnan M, La Gerche A, Kalman J, et al. Comparison of frequency of significant electrocardiographic abnormalities in endurance versus nonendurance athletes. *Am J Cardiol* 2014;113:1567-73.
68. Malhotra A, Dhutia H, Gati S, et al. Prevalence and significance of anterior T wave inversion in young white athletes and non athletes. *J Am Coll Cardiol* 2016;69:1-9.
69. Riding NR, Salah O, Sharma S, et al. ECG and morphologic adaptations in Arabic athletes: are the European Society of Cardiology's recommendations for the interpretation of the 12-lead ECG appropriate for this ethnicity? *Br J Sports Med* 2014;48:1138-43.
70. Maron BJ, Wolfson JK, Ciro E, Spirito P. Relation of electrocardiographic abnormalities and patterns of left ventricular hypertrophy identified by 2-dimensional echocardiography in patients with hypertrophic cardiomyopathy. *Am J Cardiol* 1983;51:189-94.
71. Haghjoo M, Mohammadzadeh S, Taherpour M, et al. ST-segment depression as a risk factor in hypertrophic cardiomyopathy. *Europace* 2009;11:643-9.
72. MacAlpin RN. Clinical significance of QS complexes in V1 and V2 without other electrocardiographic abnormality. *Ann Noninvasive Electrocardiol* 2004;9:39-47.
73. Bent RE, Wheeler MT, Hadley D, Froelicher V, Ashley E, Perez MV. Computerized Q wave dimensions in athletes and hypertrophic cardiomyopathy patients. *J Electrocardiol* 2015;48:362-7.
74. Riding NR, Sheikh N, Adamuz C, et al. Comparison of three current sets of electrocardiographic interpretation criteria for use in screening athletes. *Heart* 2014;101:384-90.
75. Kim JH, Baggish AL. Electrocardiographic right and left bundle branch block patterns in athletes: prevalence, pathology, and clinical significance. *J Electrocardiol* 2015;48:380-4.
76. Le VV, Wheeler MT, Mandic S, et al. Addition of the electrocardiogram to the preparticipation examination of college athletes. *Clin J Sport Med* 2010;20:98-105.
77. Aro AL, Anttonen O, Tikkanen JT, et al. Intraventricular conduction delay in a standard 12-lead electrocardiogram as a predictor of mortality in the general population. *Circ Arrhythm Electrophysiol* 2011;4:704-10.
78. Desai AD, Yaw TS, Yamazaki T, Kaykha A, Chun S, Froelicher VF. Prognostic significance of quantitative QRS duration. *Am J Med* 2006;119:600-6.
79. Dunn T, Abdelfattah R, Aggarwal S, Pickham D, Hadley D, Froelicher V. Are the QRS duration and ST depression cut-points from the Seattle criteria too conservative? *J Electrocardiol* 2015;48:395-8.
80. Xiao HB, Brecker SJ, Gibson DG. Relative effects of left ventricular mass and conduction disturbance on activation in patients with pathological left ventricular hypertrophy. *Br Heart J* 1994;71:548-53.
81. Surawicz B, Childers R, Deal BJ, et al. AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: part III: intraventricular conduction disturbances: a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society: endorsed by the International Society for Computerized Electrocardiology. *J Am Coll Cardiol* 2009;53:976-81.
82. Drezner JA, Prutkin JM, Harmon KG, et al. Cardiovascular screening in college athletes. *J Am Coll Cardiol* 2015;65:2353-5.
83. Cohen MI, Triedman JK, Cannon BC, et al. PACES/HRS expert consensus statement on the management of the asymptomatic young patient with a Wolff-Parkinson-White (WPW, ventricular preexcitation) electrocardiographic pattern: developed in partnership between the Pediatric and Congenital Electrophysiology Society (PACES) and the Heart Rhythm Society (HRS). Endorsed by the governing bodies of PACES, HRS, the American College of Cardiology Foundation (ACCF), the American Heart Association (AHA), the American Academy of Pediatrics (AAP), and the Canadian Heart Rhythm Society (CHRS). *Heart Rhythm* 2012;9:1006-24.
84. Daubert C, Ollitrault J, Descaves C, Mabo P, Ritter P, Gouffault J. Failure of the exercise test to predict the anterograde refractory period of the accessory pathway in Wolff Parkinson White syndrome. *Pacing Clin Electrophysiol* 1988;11:1130-8.
85. Klein GJ, Gulamhusein SS. Intermittent preexcitation in the Wolff-Parkinson-White syndrome. *Am J Cardiol* 1983;52:292-6.

86. Klein GJ, Bashore TM, Sellers TD, Pritchett EL, Smith WM, Gallagher JJ. Ventricular fibrillation in the Wolff-Parkinson-White syndrome. *N Engl J Med* 1979;301:1080–5.
87. Schwartz PJ, Stramba-Badiale M, Crotti L, et al. Prevalence of the congenital long-QT syndrome. *Circulation* 2009;120:1761–7.
88. Tester DJ, Ackerman MJ. Cardiomyopathic and channelopathic causes of sudden unexplained death in infants and children. *Annu Rev Med* 2009;60:69–84.
89. Eckart RE, Shry EA, Burke AP, et al. Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol* 2011;58:1254–61.
90. Meyer L, Stubbs B, Fahrenbruch C, et al. Incidence, causes, and survival trends from cardiovascular-related sudden cardiac arrest in children and young adults 0 to 35 years of age: a 30-year review. *Circulation* 2012;126:1363–72.
91. Behr E, Wood DA, Wright M, et al. Cardiologic assessment of first-degree relatives in sudden arrhythmic death syndrome. *Lancet* 2003;362:1457–9.
92. Tester DJ, Spoon DB, Valdivia HH, Makielski JC, Ackerman MJ. Targeted mutational analysis of the RyR2-encoded cardiac ryanodine receptor in sudden unexplained death: a molecular autopsy of 49 medical examiner/coroner's cases. *Mayo Clin Proc* 2004;79:1380–4.
93. Tan HL, Hofman N, van Langen IM, van der Wal AC, Wilde AA. Sudden unexplained death: heritability and diagnostic yield of cardiologic and genetic examination in surviving relatives. *Circulation* 2005;112:207–13.
94. Finocchiaro G, Papadakis M, Robertus JL, et al. Etiology of sudden death in sports: insights from a United Kingdom Regional Registry. *J Am Coll Cardiol* 2016;67:2108–15.
95. Viskin S, Rosovski U, Sands AJ, et al. Inaccurate electrocardiographic interpretation of long QT: the majority of physicians cannot recognize a long QT when they see one. *Heart Rhythm* 2005;2:569–74.
96. Postema PG, De Jong JS, Van der Bilt IA, Wilde AA. Accurate electrocardiographic assessment of the QT interval: teach the tangent. *Heart Rhythm* 2008;5:1015–8.
97. Bazett HC. An analysis of the time-relations of electrocardiograms. *Heart* 1920:353–70.
98. Malfatto G, Beria G, Sala S, Bonazzi O, Schwartz PJ. Quantitative analysis of T wave abnormalities and their prognostic implications in the idiopathic long QT syndrome. *J Am Coll Cardiol* 1994;23:296–301.
99. Ackerman MJ, Zipes DP, Kovacs RJ, Maron BJ. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: Task Force 10: The cardiac channelopathies: a scientific statement from the American Heart Association and American College of Cardiology. *J Am Coll Cardiol* 2015;66:2424–8.
100. Dhutia H, Malhotra A, Parpia S, et al. The prevalence and significance of a short QT interval in 18 825 low-risk individuals including athletes. *Br J Sports Med* 2016;50:124–9.
101. Basavarajiah S, Wilson M, Whyte G, Shah A, Behr E, Sharma S. Prevalence and significance of an isolated long QT interval in elite athletes. *Eur Heart J* 2007;28:2944–9.
102. Goldenberg I, Moss AJ, Peterson DR, et al. Risk factors for aborted cardiac arrest and sudden cardiac death in children with the congenital long-QT syndrome. *Circulation* 2008;117:2184–91.
103. Gollob MH, Redpath CJ, Roberts JD. The short QT syndrome: proposed diagnostic criteria. *J Am Coll Cardiol* 2011;57:802–12.
104. Schwartz PJ, Moss AJ, Vincent GM, Crampton RS. Diagnostic criteria for the long QT syndrome. An update. *Circulation* 1993;88:782–4.
105. Schwartz PJ, Crotti L. QTC behavior during exercise and genetic testing for the long-QT syndrome. *Circulation* 2011;124:2181–4.
106. Ackerman MJ, Priori SG, Willems S, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm* 2011;8:1308–39.
107. Wilde AA, Antzelevitch C, Borggrefe M, et al. Proposed diagnostic criteria for the Brugada syndrome: consensus report. *Circulation* 2002;106:2514–9.
108. Bayes de Luna A, Brugada J, Baranchuk A, et al. Current electrocardiographic criteria for diagnosis of Brugada pattern: a consensus report. *J Electrocardiol* 2012;45:433–42.
109. Priori SG, Wilde AA, Horie M, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm* 2013;10:1932–63.
110. Zorzi A, Leoni L, Di Paolo FM, et al. Differential diagnosis between early repolarization of athlete's heart and coved-type Brugada electrocardiogram. *Am J Cardiol* 2015;115:529–32.
111. Verdile L, Maron BJ, Pelliccia A, Spataro A, Santini M, Biffi A. Clinical significance of exercise-induced ventricular tachyarrhythmias in trained athletes without cardiovascular abnormalities. *Heart Rhythm* 2015;12:78–85.
112. Biffi A, Pelliccia A, Verdile L, Fernando F, Spataro A, Caselli S, Santini M, Maron BJ. Long-term clinical significance of frequent and complex ventricular tachyarrhythmias in trained athletes. *J Am Coll Cardiol* 2002;40:446–52.
113. Novak J, Zorzi A, Castelletti S, et al. Electrocardiographic differentiation of idiopathic right ventricular outflow tract ectopy from early arrhythmogenic right ventricular cardiomyopathy. *Europace* 2016 Jun 23 [E-pub ahead of print].
114. Corrado D, Basso C, Leoni L, et al. Three-dimensional electroanatomical voltage mapping and histologic evaluation of myocardial substrate in right ventricular outflow tract tachycardia. *J Am Coll Cardiol* 2008;51:731–9.
115. Heidbuchel H, Hoogsteen J, Fagard R, et al. High prevalence of right ventricular involvement in endurance athletes with ventricular arrhythmias. Role of an electrophysiologic study in risk stratification. *Eur Heart J* 2003;24:1473–80.
116. Biffi A, Maron BJ, Verdile L, et al. Impact of physical deconditioning on ventricular tachyarrhythmias in trained athletes. *J Am Coll Cardiol* 2004;44:1053–8.
117. Delise P, Lanari E, Sitta N, Centa M, Allocca G, Biffi A. Influence of training on the number and complexity of frequent VPBs in healthy athletes. *J Cardiovasc Med (Hagerstown)* 2011;12:157–61 [PMC [10.2459/JCM.0b013e32834102ea] [21139509].
118. Delise P, Sitta N, Lanari E, et al. Long-term effect of continuing sports activity in competitive athletes with frequent ventricular premature complexes and apparently normal heart. *Am J Cardiol* 2013;112:1396–402.
119. Anderson JL, Halperin JL, Albert NM, et al. Management of patients with atrial fibrillation (compilation of 2006 ACCF/AHA/ESC and 2011 ACCF/AHA/HRS recommendations): a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;61:1935–44.
120. Aagaard P, Sahlen A, Braunschweig F. Performance trends and cardiac biomarkers in a 30-km cross-country race, 1993–2007. *Med Sci Sports Exerc* 2012;44:894–9.
121. Sahlen A, Gustafsson TP, Svensson JE, et al. Predisposing factors and consequences of elevated biomarker levels in long-distance runners aged >or=55 years. *Am J Cardiol* 2009;104:1434–40.
122. Rose G, Baxter PJ, Reid DD, McCartney P. Prevalence and prognosis of electrocardiographic findings in middle-aged men. *Br Heart J* 1978;40:636–43.
123. Chou R, Arora B, Dana T, Fu R, Walker M, Humphrey L. Screening asymptomatic adults with resting or exercise electrocardiography: a review of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2011;155:375–85.
124. Daviglus ML, Liao Y, Greenland P, et al. Association of nonspecific minor ST-T abnormalities with cardiovascular mortality: the Chicago Western Electric Study. *Jama* 1999;281:530–6.
125. Borjesson M, Urhausen A, Koudi E, et al. Cardiovascular evaluation of middle-aged/senior individuals engaged in leisure-time sport activities: position stand from the sections of exercise physiology and sports cardiology of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2011;18:446–58.
126. Maron BJ, Araujo CG, Thompson PD, et al. Recommendations for preparticipation screening and the assessment of cardiovascular disease in masters athletes: an advisory for healthcare professionals from the working groups of the World Heart Federation, the International Federation of Sports Medicine, and the American Heart Association Committee on Exercise, Cardiac Rehabilitation, and Prevention. *Circulation* 2001;103:327–34.

KEY WORDS athlete, electrocardiogram

APPENDIX For supplemental material containing an extended account of the document and additional illustrative ECGs, please see the online version of this article.